

*New Idea*

Vacuums of the sea: Ecological function of large coral reef benthic scavengers in suppressing crown-of-thorns starfish (COTS) outbreaks

Mohsen Kayal and Hunter S. Lenihan

Mohsen Kayal (mohsen.kayal@ird.fr), ENTROPIE, IRD, Université de la Réunion, CNRS, IFREMER, Université de la Nouvelle-Calédonie, Noumea, Kanaky New Caledonia and Laboratoire d'Excellence CORAIL, Paris, France

Hunter S. Lenihan (hlenihan@ucsb.edu), Bren School of Environmental Science and Management, University of California, Santa Barbara, USA

Abstract

Despite their drastic impacts on coral reefs, outbreaks of the coral-feeding seastar crown-of-thorns starfish (COTS), *Acanthaster*, have remained a scientific enigma. Significant efforts in coral reef conservation science have been dedicated to identifying natural predators able to exert demographic control on COTS and prevent population outbreaks. These efforts are motivated by empirical evidence showing that reefs within marine protected areas are less prone to COTS outbreaks than reefs open to fishing where potential COTS predators have been reduced or removed functionally from food webs. Research findings point towards COTS' early life-stages as a major demographic bottleneck for COTS populations, with various reef fish and benthic organisms identified as predators of the seastar. Yet, no species or species groups have been clearly identified as exerting enough top-down control to influence COTS population increases or prevent outbreaks. We report the benthic scavenging behavior of eagle rays (family *Myliobatidae*), a large-bodied predator, feeding in coral rubble fields of Kanaky New Caledonia, critical habitats where juvenile COTS find refuge and food and accumulate to produce population outbreaks. We argue that with their effective substrate-sucking feeding behavior, similar to vacuums of the sea, eagle rays may be a hitherto unidentified predator able to exert significant control on COTS populations. Eagle rays and other large benthic scavengers were previously neglected in the search for major COTS predators. Relatively little existing data

show that eagle ray populations in Kanaky New Caledonia's lagoon are more abundant inside than outside marine protected areas, which concurs with the hypothesis that they could be responsible for the mitigation of COTS outbreaks as reported within reserves. We advocate for further investigations on the role of eagle rays and other large benthic scavengers in controlling COTS outbreaks, and the importance of preserving the unique ecological function of sea vacuums for coral reef conservation.

Keywords: Coral reef; *Acanthaster*; Seastar; Predator outbreak; Eagle ray; Nature-based solution.

Unresolved mystery surrounding COTS outbreaks

Outbreaks of crown-of-thorns starfish (COTS, *Acanthaster* spp.) cause mass coral mortality and are a clear threat to coral reef conservation in an era of global coral decline (Kayal et al. 2012, Pratchett et al. 2017, Condie et al. 2021). There has been a long-running debate about the mechanisms that control COTS outbreaks, which, in part, remain a scientific enigma. Two predominant hypotheses about the mechanisms exerting demographic bottlenecks for COTS populations have been proposed, both of which are supported by a relatively large body of empirical evidence (Pratchett et

al. 2014). The bottom-up “nutrient enrichment hypothesis” emphasizes the importance of nutrient availability for phytoplankton blooms upon which COTS larvae feed and rely during their transition from pelagic to benthic stages: with eutrophication of reef environments, COTS populations increase due to increased recruitment and early life stage survivorship (Fabricius et al. 2010, Wooldridge and Brodie 2015, Matthews et al. 2020). The top-down “predation release hypothesis” emphasizes the role of species feeding upon the seastar in regulating their populations, a mechanism altered when reef fisheries reduce local abundances of COTS predators (Cowan et al. 2017). Recently, Deaker et al. (2020) discovered a third mechanism by showing that COTS is able to remain in an “ecologically dormant” state of herbivorous juvenile hiding in dead coral rubble, enabling the build-up of cryptic outbreak populations over years, before shifting to corallivory and full-grown adults once preferred coral prey is available (Wilmes et al. 2020a, Deaker and Byrne 2022, Neil et al. 2022). Quantitative evidence indicates that COTS outbreaks are more frequent and intense outside of marine protected areas than in reef areas protected from fishing, thereby supporting the predation release hypothesis (Dulvy et al. 2004, Sweatman 2008, McCook et al. 2010, Mellin et al. 2016, Vanhatalo et al. 2017, Westcott et al. 2020, Kroon et al. 2021). Nevertheless, the evidence to support the

predation release hypothesis so far remains predominantly correlative and circumstantial, and there is need for improved understanding of the mechanisms through which protection from fishing benefits control of COTS populations. Specifically, identifying species exerting significant control on COTS populations can help define targeted management actions that support coral reef resilience.

Our hypothesis

We propose the hypothesis that foraging by eagle rays (family *Myliobatidae*), large-bodied bottom scavengers feeding in coral reef rubble fields, can exert significant top-down control of COTS juvenile life-stages, thereby playing a key role in regulating COTS outbreaks. Our hypothesis is based on observations of active foraging of the reef substrate in Kanaky New Caledonia by an eagle ray, moving coral rubble around and sucking up potential prey hidden in the substrate and around live corals (Figure 1). Because coral rubble fields constitute predominant habitats for COTS juvenile stages (Zann et al. 1987, Wilmes et al. 2020b), we argue that such feeding behavior, acting as a vacuum of the sea substrate, may constitute a key ecological function regulating marine benthic invertebrate populations, including COTS whose outbreaks can devastate corals at large spatial scales.



Figure 1. Photograph of an eagle ray (family *Myliobatidae*) foraging the shallow reef substrate in Kanaky New Caledonia. The individual was observed moving coral rubbles and sucking up potential prey from the reef substrate, a suction feeding behavior comparable to a vacuum of the sea. We believe that this ecological function may be critical for regulating invertebrate populations hiding in the substrate, such as juveniles of the coral predator crown-of-thorns starfish. See video at Kayal (2024).

Support for our hypothesis

Recent research efforts have focused on identifying COTS predators in an endeavor to improve coral reef management through the regulation of COTS outbreaks exerted by their natural predators. More than 100 species of reef fish and invertebrates have been identified as preying upon the seastar at different life stages and health conditions (Cowan et al. 2017). However, predation on early COTS life stages remains largely under-characterized, while seastar survival in the juvenile stage—that is when the small, cryptic, and coralline feeding seastars are hiding within coral rubble habitats—is considered a major demographic bottleneck for COTS populations (Wilmes et al. 2018, 2020b). In an extensive review of the scientific literature, Cowan et al. (2017) concluded that predation on newly settled starfish is predominantly exerted by benthic invertebrates associated with coral rubble fields, a hypothesis currently under intensive investigation (Desbiens et al. 2023). However, research on putative predators of juvenile COTS have hitherto largely dismissed the importance of large generalist reef predators that are able to effectively forage through coral rubble over expansive reef areas, like eagle rays. Eagle rays, like other large rays, are intensive bottom-feeders that effectively “vacuum” up invertebrate prey hidden in marine benthic substrate (Thrush et al. 1991, Hines et al. 1997, Ajemian et al. 2018, Figure 1). A large body of research conducted in the 1980's solidified how foraging by bat rays in temperate latitude regions (Hulberg and Oliver 1980, Van Blaricom 1982), and sting rays in tropical latitudes (Thistle 1981), substantially influence soft-sediment invertebrate assemblages, either directly through predation or indirectly through physical disturbance (see Lenihan and Micheli 2001 for a review). Our observation of an eagle ray foraging in coral rubble suggests that they may be a key predator able to control COTS juvenile abundance and reduce outbreaks. Indeed, coral rubble provides important recruitment substrate, refuge, and coralline-algae food for juvenile COTS (Zann et al. 1987, Wilmes et al. 2018, 2020a), and is where seastar density can build up over years leading to outbreaks (Deaker et al. 2020).

Juvenile COTS buried in coral rubble may not have many large predators (Cowan et al. 2017). Species acting as sea vacuums in rubble fields, like eagle rays, may play a unique role in controlling COTS density build-ups and the frequency and intensity of outbreaks, which typically originate in habitats dominated by coral rubble (Zann et al. 1987, Johnson et al. 1991, Kayal et al. 2012, Wilmes et al. 2020b). Relatively healthy eagle ray populations may explain the relatively constrained COTS outbreaks observed in Kanaky New Caledonia's lagoon, as compared with the neighboring regions Australia and Vanuatu, where higher fishing and shark-net impacts on

rays and higher frequency and intensity of COTS outbreaks are reported (Lynch et al. 2010, Schluessel et al. 2010, Harry et al. 2011, Sumpton et al. 2011, Adjerdoud et al. 2018, Sporčić et al. 2018, Laran et al. 2024). Although data limitations currently prevent quantitatively testing our hypothesis, recent investigation in Kanaky New Caledonia indicates eagle ray populations were three times more abundant inside than outside of marine reserves (Heudier et al. 2023), supporting the hypothesis that they could be responsible for the mitigation of COTS outbreaks in protected areas, as reported in other regions (Dulvy et al. 2004, Sweatman 2008, McCook et al. 2010, Mellin et al. 2016, Westcott et al. 2020, Kroon et al. 2021). Other hitherto overlooked benthic scavenger species potentially exerting significant predatory control on juvenile COTS hidden in coral rubble and with populations severely impacted by fisheries include sea cucumbers (Holothuria; Anderson et al. 2011, Purcell et al. 2013, Pierrat et al. 2022, Ahmed et al. 2023), although this is beyond the scope of the present study.

Conclusions and prospective research

Megafauna often play important roles in marine ecosystems, although their ecological functions are not always known or fully recognized, and their populations are often significantly reduced by human activities (McCauley et al. 2015, Pimiento et al. 2020). Our observations shed light on the potentially unique ecological function of eagle rays for safeguarding coral reefs from COTS outbreaks. While we currently do not possess the necessary information to test this hypothesis, future investigations may include eagle ray behavior and diet analyses to confirm or reject predation on juvenile COTS (Ajemian et al. 2012, Leray et al. 2013), and joint analysis of eagle ray and COTS populations to establish predator-prey regulation and a potential mitigation of outbreaks (Kayal et al. 2012). While diver operated COTS control efforts are costly but often considered as the best approach to reduce the impacts of their outbreaks (Westcott et al. 2020, Condie et al. 2021, Castro-Sanguino et al. 2023), coral reef resilience may significantly benefit from an increased protection of eagle ray populations and their important ecological function as vacuums of the reef.

Acknowledgements

This study is part of the Track Changes (2025) program and was supported by funding from the French Laboratory of Excellence CORAIL through a starting grant, and the French Ministry of Foreign Affairs through the regional cooperation grant Fonds Pacifique. A first version of this manuscript was made available as a preprint (Kayal and Lenihan 2024).

Referees

Benjamin Mos – University of Queensland
b.mos@uq.edu.au

Morgan Pratchett – James Cook University
morgan.pratchett@jcu.edu.au

References

- Adjeroud, M., Kayal, M., Peignon, C., Juncker, M., Mills, S.C., Beldade, R., et al. 2018. Ephemeral and localized outbreaks of the coral predator *Acanthaster* cf. *solaris* in the southwestern lagoon of New Caledonia. *Zoological Studies* 57: 4. [CrossRef](#)
- Ahmed, Q., Ali, Q.M., Bat, L., Öztekin, A., Ghory, F. S., Shaikh, I., et al. 2023. Gut content analysis in *Holothuria leucospilota* and *Holothuria cinerascens* (Echinodermata: Holothuroidea: Holothuriidae) From Karachi Coast. *Journal of Materials and Environmental Science* 14: 31–40. https://www.imatrenvironsci.com/Document/vol14/vol14_N1/JMES-2023-14003-Ahmed.pdf
- Ajemian, M.J., Powers, S.P. and T.J.T. Murdoch. 2012. Estimating the potential impacts of large mesopredators on benthic resources: integrative assessment of spotted eagle ray foraging ecology in Bermuda. *PLOS ONE* 7: e40227. [CrossRef](#)
- Anderson, S.C., Flemming, J.M., Watson, R. and H.K. Lotze. 2011. Serial exploitation of global sea cucumber fisheries. *Fish and Fisheries* 12: 317–339. [CrossRef](#)
- Castro-Sanguino, C., Bozec, Y.M., Condie, S.A., Fletcher, C.S., Hock, K., Roelfsema, C., et al. 2023. Control efforts of crown-of-thorns starfish outbreaks to limit future coral decline across the Great Barrier Reef. *Ecosphere* 14: e4580. [CrossRef](#)
- Condie, S.A., Anthony, K.R., Babcock, R.C., Baird, M.E., Beeden, R., Fletcher, C.S., et al. 2021. Large-scale interventions may delay decline of the Great Barrier Reef. *Royal Society Open Science* 8: 201296. [CrossRef](#)
- Cowan, Z.L., Pratchett, M., Messmer, V. and S. Ling. 2017. Known predators of crown-of-thorns starfish (*Acanthaster* spp.) and their role in mitigating, if not preventing, population outbreaks. *Diversity* 9: 7. [CrossRef](#)
- Deaker, D.J., Agüera, A., Lin, H.A., Lawson, C., Budden, C., Dworjanyn, S.A., et al. 2020. The hidden army: corallivorous crown-of-thorns seastars can spend years as herbivorous juveniles. *Biology Letters* 16: 20190849. [CrossRef](#)
- Deaker, D.J. and M. Byrne. 2022. Crown of thorns starfish life-history traits contribute to outbreaks, a continuing concern for coral reefs. *Emerging Topics in Life Sciences* 6: 67–79. [CrossRef](#)
- Desbiens, A.A., Mumby, P.J., Dworjanyn, S., Plagányi, É.E., Uthicke, S. and K. Wolfe. 2023. Novel rubble-dwelling predators of herbivorous juvenile crown-of-thorns starfish (*Acanthaster* sp.). *Coral Reefs* 42: 579–591. [CrossRef](#)
- Dulvy, N.K., Freckleton, R.P. and N.V. Polunin. 2004. Coral reef cascades and the indirect effects of predator removal by exploitation. *Ecology Letters* 7: 410–416. [CrossRef](#)
- Fabricius, K.E., Okaji, K. and G. De'ath. 2010. Three lines of evidence to link outbreaks of the crown-of-thorns seastar *Acanthaster planci* to the release of larval food limitation. *Coral Reefs* 29: 593–605. [CrossRef](#)
- Harry, A.V., Tobin, A.J., Simpfendorfer, C.A., Welch, D.J., Mapleston, A., White, J., et al. 2011. Evaluating catch and mitigating risk in a multispecies, tropical, inshore shark fishery within the Great Barrier Reef World Heritage Area. *Marine and Freshwater Research* 62: 710–721. [CrossRef](#)
- Heudier, M., Mouillot, D. and L. Mannocci. 2023. Assessing the effects of coral reef habitat and marine protected areas on threatened megafauna using aerial surveys. *Aquatic Conservation: Marine and Freshwater Ecosystems* 33: 286–297. [CrossRef](#)
- Hines, A.H., Whitlatch, R.B., Thrush, S.F., Hewitt, J.E., Cummings, V.J., Dayton, P., et al. 1997. Nonlinear foraging response of a large marine predator to benthic prey: eagle ray pits and bivalves in a New Zealand sandflat. *Journal of Experimental Marine Biology and Ecology* 216: 191–210. [CrossRef](#)
- Hulberg, L.W. and J.S. Oliver. 1980. Caging manipulations in marine soft-bottom communities: importance of animal interactions or sedimentary habitat modifications. *Canadian Journal of Fisheries and Aquatic Sciences* 37: 1130–1139. [CrossRef](#)
- Johnson, C.R., Sutton, D.C., Olson, R.R. and R. Giddins. 1991. Settlement of crown-of-thorns starfish: role of bacteria on surfaces of coralline algae and a hypothesis for deepwater recruitment. *Marine Ecology Progress Series* 71: 143–162. <https://www.int-res.com/articles/meps/71/m071p143.pdf>
- Kayal, M. 2024. Eagle ray feeding on a shallow coral reef in Kanaky New Caledonia. Aerial video footage, 2 min 45 sec. <https://youtu.be/sxdbiZJmX1Y>
- Kayal, M. and H.S. Lenihan. 2024. Vacuums of the sea: ecological function of large coral reef benthic scavengers in suppressing crown-of-thorns starfish (COTS) outbreaks. *Zenodo*. [CrossRef](#)
- Kayal, M., Vercelloni, J., Lison de Loma, T., Bosserelle, P., Chancerelle, Y., Geoffroy, S., et al. 2012. Predator crown-of-thorns starfish (*Acanthaster planci*) outbreak, mass mortality of corals, and cascading effects on reef fish and benthic communities. *PLOS ONE*: e47363. [CrossRef](#)

- Kroon, F.J., Barneche, D.R. and M.J. Emslie. 2021. Fish predators control outbreaks of Crown-of-Thorns Starfish. *Nature Communications* 12: 6986. [CrossRef](#)
- Laran, S., Van Canneyt, O., Dorémus, G., Garrigue, C., Berr, T., Bourgoigne, H., et al. 2023. Who lives in the open sea? Distribution and densities of surfacing marine megafauna in three subregions of the South Pacific (New Caledonia, Wallis and Futuna, and French Polynesia). *Pacific Conservation Biology* 30: PC23023. [CrossRef](#)
- Lenihan, H.S. and F. Micheli. 2001. Soft sediment communities. Pages 253–287 in Bertness, M., Hay, M.E., and S.D. Gaines, editors. *Marine Community Ecology*. Sinauer Associates.
- Leray, M., Yang, J.Y., Meyer, C.P., Mills, S.C., Agudelo, N., Ranwez, V., et al. 2013. A new versatile primer set targeting a short fragment of the mitochondrial COI region for metabarcoding metazoan diversity: application for characterizing coral reef fish gut contents. *Frontiers in Zoology* 10: 1–14. [CrossRef](#)
- Lynch, A.M.J., Sutton, S.G. and C.A. Simpfendorfer. 2010. Implications of recreational fishing for elasmobranch conservation in the Great Barrier Reef Marine Park. *Aquatic Conservation: Marine and Freshwater Ecosystems* 20: 312–318. [CrossRef](#)
- Matthews, S.A., Mellin, C. and M.S. Pratchett. 2020. Larval connectivity and water quality explain spatial distribution of crown-of-thorns starfish outbreaks across the Great Barrier Reef. *Advances in Marine Biology* 87: 223–258. [CrossRef](#)
- McCauley, D.J., Pinsky, M.L., Palumbi, S.R., Estes, J.A., Joyce, F.H. and R.R. Warner. 2015. Marine defaunation: animal loss in the global ocean. *Science* 347: 1255641. [CrossRef](#)
- McCook, L.J., Ayling, T., Cappo, M., Choat, J.H., Evans, R.D., De Freitas, D.M., et al. 2010. Adaptive management of the Great Barrier Reef: a globally significant demonstration of the benefits of networks of marine reserves. *Proceedings of the National Academy of Sciences* 107: 18278–18285. [CrossRef](#)
- Mellin, C., MacNeil, A.M., Cheal, A.J., Emslie, M.J. and J.M. Caley. 2016. Marine protected areas increase resilience among coral reef communities. *Ecology Letters* 19: 629–637. [CrossRef](#)
- Neil, R.C., Gomez Cabrera, M. and S. Uthicke. 2022. Juvenile age and available coral species modulate transition probability from herbivory to corallivory in *Acanthaster cf. solaris* (Crown-of-Thorns Seastar). *Coral Reefs* 41: 843–848. [CrossRef](#)
- Pierrat, J., Bédier, A., Eeckhaut, I., Magalon, H. and P. Frouin. 2022. Sophistication in a seemingly simple creature: a review of wild holothurian nutrition in marine ecosystems. *Biological Reviews* 97: 273–298. [CrossRef](#)
- Pimiento, C., Leprieur, F., Silvestro, D., Lefcheck, J.S., Albouy, C., Rasher, D.B., et al. 2020. Functional diversity of marine megafauna in the Anthropocene. *Science Advances* 6: eaay7650. [CrossRef](#)
- Pratchett, M.S., Caballes, C.F., Rivera-Posada, J.A. and H.P. Sweatman. 2014. Limits to Understanding and Managing Outbreaks of Crown-of-Thorns Starfish (*Acanthaster* spp.). *Oceanography and Marine Biology: an annual review* 52: 133–244. CRC Press.
- Pratchett, M.S., Caballes, C.F., Wilmes, J.C., Matthews, S., Mellin, C., Sweatman, H.P., et al. 2017. Thirty years of research on crown-of-thorns starfish (1986–2016): scientific advances and emerging opportunities. *Diversity* 9: 41. [CrossRef](#)
- Purcell, S.W., Mercier, A., Conand, C., Hamel, J.F., Toral-Granda, M.V., Lovatelli, A., et al. 2013. Sea cucumber fisheries: global analysis of stocks, management measures and drivers of overfishing. *Fish and Fisheries* 14: 34–59. [CrossRef](#)
- Schluessel, V., Bennett, M.B. and S.P. Collin. 2010. Diet and reproduction in the white-spotted eagle ray *Aetobatus narinari* from Queensland, Australia and the Penghu Islands, Taiwan. *Marine and Freshwater Research* 61: 1278–1289. [CrossRef](#)
- Sporcic, M., Hobday, A., Bulman, C. and M. Fuller. 2018. Ecological risk assessment for the effects of fishing: Eastern tuna & billfish fishery: Longline Sub-fishery, data to 2015. Report for the Australian Fisheries Management Authority.
- Sumpton, W.D., Taylor, S.M., Gribble, N.A., McPherson, G. and T. Ham. 2011. Gear selectivity of large-mesh nets and drumlines used to catch sharks in the Queensland Shark Control Program. *African Journal of Marine Science* 33: 37–43. [CrossRef](#)
- Sweatman, H. 2008. No-take reserves protect coral reefs from predatory starfish. *Current Biology* 18: R598–9. [CrossRef](#)
- Thistle, D. 1981. Natural physical disturbances and communities of marine soft bottoms. *Marine Ecology Progress Series* 6: 223–228. <https://www.int-res.com/articles/meps/6/m006p223.pdf>
- Thrush, S.F., Pridmore, R.D., Hewitt, J.E. and V.J. Cummings. 1991. Impact of ray feeding disturbances on sandflat macrobenthos: do communities dominated by polychaetes or shellfish respond differently? *Marine Ecology Progress Series* 69: 245–252. <https://www.int-res.com/articles/meps/69/m069p245.pdf>
- Track Changes 2025. Coastal Marine Ecosystem Transformation in the Face of Environmental Changes. Accessed 12 February 2025 <https://umr-entropie.ird.nc/index.php/home-en/home/research/portfolio/ongoing-projects/projet-track-changes>
- VanBlaricom, G.R. 1982. Experimental analyses of structural regulation in a marine sand community exposed to oceanic swell. *Ecological Monographs* 52: 283–305. [CrossRef](#)

- Vanhatalo, J., Hosack, G.R. and H. Sweatman. 2017. Spatiotemporal modelling of crown-of-thorns starfish outbreaks on the Great Barrier Reef to inform control strategies. *Journal of Applied Ecology* 54: 188–197. [CrossRef](#)
- Westcott, D.A., Fletcher, C.S., Kroon, F.J., Babcock, R.C., Plagányi, E.E., Pratchett, M.S., et al. 2020. Relative efficacy of three approaches to mitigate Crown-of-Thorns Starfish outbreaks on Australia's Great Barrier Reef. *Scientific Reports* 10: 12594. [CrossRef](#)
- Wilmes, J.C., Caballes, C.F., Cowan, Z.L., Hoey, A.S., Lang, B.J., Messmer, V. and M.S. Pratchett. 2018. Contributions of pre-versus post-settlement processes to fluctuating abundance of crown-of-thorns starfishes (*Acanthaster* spp.). *Marine Pollution Bulletin* 135: 332–345. [CrossRef](#)
- Wilmes, J.C., Hoey, A.S. and M.S. Pratchett. 2020a. Contrasting size and fate of juvenile crown-of-thorns starfish linked to ontogenetic diet shifts. *Proceedings of the Royal Society B* 287: 20201052. [CrossRef](#)
- Wilmes, J.C., Schultz, D.J., Hoey, A.S., Messmer, V. and M.S. Pratchett. 2020b. Habitat associations of settlement-stage crown-of-thorns starfish on Australia's Great Barrier Reef. *Coral Reefs* 39: 1163–1174. [CrossRef](#)
- Wooldridge, S.A. and J.E. Brodie. 2015. Environmental triggers for primary outbreaks of crown-of-thorns starfish on the Great Barrier Reef, Australia. *Marine Pollution Bulletin* 101: 805–815. [CrossRef](#)
- Zann L., Brodie J., Berryman C. and M. Naqasima. 1987. Recruitment, ecology, growth and behavior of juvenile *Acanthaster planci* (L.) (Echinodermata: Asteroidea). *Bulletin of Marine Science* 41: 561–575.

Response to referee

We agree with Pratchett (2025) that dedicated studies are necessary before confirming our hypothesis that eagle rays exert regulatory pressure on juvenile populations of the coral-eating seastar crown-of-thorns starfish (COTS). As stated in our original manuscript, and indicated by the journal name—*Ideas in Ecology and Evolution*—and category of article we have proposed, our hypothesis remains predominantly a *new idea* that requires further investigation. The discussion platform provided by this journal enables further debating the mechanisms underlying our hypothesis with colleagues working on the topic, and identifying ways to test them in future research. There are predominantly two lines of evidence, one mechanistic and the other correlative, pointing towards the potential role of eagle rays in regulating COTS juvenile abundance in coral reef rubble fields.

Mechanistic line of evidence: eagle rays are large-bodied generalist vacuum feeders able to suck up large quantity of prey from coral reef rubble, which is the predominant habitat in which juvenile COTS find refuge to build up outbreaking populations.

We agree that studies on the feeding behavior of eagle rays are necessary to confirm the commonness of our observation that eagle rays regularly prey in coral reef rubble fields, and that their feeding activity has indeed regulatory effects on COTS populations. Until such knowledge is available there is no reason to believe that our observation is an exclusive behavior (i.e. a one-time event, or that of a deviant individual), or that eagle rays' vacuum feeding in coral reef rubble would spare juvenile COTS. The eagle ray in our video is seen displacing a relatively large piece of coral rubble before scrutinizing the area and overturned rubble piece with its mouth, seemingly sucking up uncovered small prey living in the interstices (see Kayal 2024 video at 1:00-1:20). It seems probable that such feeding behavior would exert strong indiscriminate predation on meio- and macro-benthic invertebrates within the size range of a millimeter to few centimeters, including juvenile COTS (Hines et al. 1997, Ajemian et al. 2012, Deaker et al. 2020, Wilmes et al. 2020a,b). We are not stipulating that eagle rays specifically search for and actively target juvenile COTS hidden in rubble fields in a way to fully suppress their populations even at low densities, but rather that repeated eagle ray vacuum feeding in rubble fields may result in keeping COTS populations in check.

As highlighted by Pratchett (2025), few studies have reported echinoderms in eagle ray diet (Capapé 1976, Gray et al. 1997, Schluessel et al. 2010, Ajemian et al. 2012, Serrano-Flores et al. 2018), and to our knowledge there is currently no evidence that eagle rays do consume COTS juveniles. However, there is no evidence that eagle rays avoid preying upon juvenile COTS when vacuum feeding in coral reef rubble. Studies describing stomach content of eagle rays and affiliates report a diversified diet, with a large predominance of mollusks and few echinoderms, sometimes including seastars (Capapé 1976), in proportions predominantly reflecting local prey composition (Gray et al. 1997, Jardas et al. 2004, Yamaguchi et al. 2005, Schluessel et al. 2010, Serrano-Flores et al. 2018, Cahill et al. 2023). Given the described generalist predator behavior with a flexible diet adapting to various environments, one can anticipate that eagle rays' indiscriminate vacuum feeding in reef substrate infested with COTS juveniles would result in high proportions in stomach contents. Among regions where COTS is present and exhibits population outbreaks, Schluessel et al. (2010) identified only small amounts of ophiuroids (another class of echinoids) in eagle ray stomachs collected in February 2007 from Australia's Heron Island, southern Great Barrier Reef,

which may have not corresponded to a time and place where high juvenile COTS densities were present in the substrate. Furthermore, visual identification of juvenile COTS in stomach contents may be challenging, as individuals are small (a millimeter to few centimeters) and lack hard structures that could facilitate identification of their remains (Deaker et al. 2020, Wilmes et al. 2020a,b, Cahill et al. 2023). As reminded by Pratchett (2025), molecular tools can complement the visual identification of prey in eagle ray stomach.

As pointed out by the referees, future research on the feeding behavior and efficiency of eagle rays may help confirm predation on juvenile COTS, provided that eagle rays and coral rubble hosting COTS juveniles are at hand, which is logistically difficult. As large coastal species inhabiting shallow lagoons and bays, eagle rays and affiliates are particularly vulnerable to fisheries and frequently caught by long- and drum-lines, nets, traps, and trawlers (Gray et al. 1997, Jardas et al. 2004, Lynch et al. 2010, Schluessel et al. 2010, Sumpton et al. 2011, Serrano-Flores et al. 2018). Therefore working with fishing communities can facilitate the acquisition of live and dead specimen for stomach content analysis and

experiments. For example, patches of coral rubble hosting juvenile COTS, as identified in situ by Wilmes et al. (2020b) or cured in aquarium by Deaker et al. (2020), can be exposed to eagle ray feeding to estimate the species' efficiency in regulating juvenile COTS populations. Such enclosure-exclosure experiments can be deployed in natural eagle ray feeding grounds as identified in our video, or in aquarium (Hines et al. 1997, Kayal et al. 2011, Ajemian et al. 2012, Flowers et al. 2021). The studies would also offer the opportunity to explore the mechanisms through which eagle ray feeding impacts the juvenile seastars at different development stages through direct predation (swallowing individuals) versus indirectly through injuries from suction and coral rubble movement, or exposure to other predators (Flowers et al. 2021). In our video, a dozen fish of various sizes are seen circling around the eagle ray, seemingly to benefit from prey exposed from the feeding activity, particularly fish from the family Labridae (probably *Thalassoma janssenii*, *Thalassoma hardwicke*, and/or *Hemigymnus fasciatus*) which are commonly observed with such behavior (Figure 2; Sabino et al. 2017, Bonham and Silbiger 2024).



Figure 2 Portions of the video frames where fish following the eagle ray can be seen, seemingly to benefit from prey exposed by the feeding activity. Thanks to their movement, the fish are easier to spot in the video, which can be watched at slow motion and high quality at Kayal (2024).

Correlative line of evidence: eagle rays are more abundant inside than outside marine protected areas, which may explain why COTS outbreaks are less preeminent in reef protected from fishing.

There is currently limited data available in our study system of Kanaky New Caledonia to confirm that eagle ray abundances are higher inside than outside protected areas at a larger scale than that demonstrated by Heudier et al. (2023), and to test to which degree ray abundances may negatively correlate with COTS populations. What is known from the literature is that, like other megafauna inhabiting coastal waters, eagle rays are highly vulnerable to human activities and declining, particularly due to targeted and incidental catch and habitat degradation (Dulvy et al. 2021, Glaus et al. 2024). Eagle rays and affiliates are frequently caught by fishing devices and shark-nets (Gray et al. 1997, Jardas et al. 2004, Serrano-Flores et al. 2018, Broadhurst and Cullis 2020), including in tropical coral reef systems such as the Australian Great Barrier Reef (Lynch et al. 2010, Schluessel et al. 2010, Harry et al. 2011, Sumpton et al. 2011, Sporic et al. 2018). In absence of further knowledge, it seems realistic to stipulate that eagle ray populations in the Great Barrier Reef may exhibit similar patterns than those observed in the Kanaky New Caledonian lagoon where densities were significantly (3-folds) higher inside than outside protected areas (Heudier et al. 2023). This concurs with the hypothesis that eagle rays may exert some level of regulation on COTS populations as found in reefs protected from fishing (Dulvy et al. 2004, Sweatman 2008, McCook et al. 2010, Mellin et al. 2016, Vanhatalo et al. 2017, Westcott et al. 2020, Kroon et al. 2021).

Correlative studies contribute greatly to characterizing predator-prey interactions and trophic cascades that trickle down across ecosystem compartments (Myers et al. 2007, Heithaus et al. 2008, Kayal et al. 2012, Flowers et al. 2021). Where spatio-temporal data on both eagle ray and COTS is available, such analysis can help testing to which degree the abundances of the two species are (negatively) correlated. As eagle rays are hard to approach in nature and constitute a diversified species group with potentially various life histories to investigate, data acquisitions can be accelerated using aerial and underwater imagery and telemetry, citizen science, environmental-DNA, and artificial intelligence (Tagliafico et al. 2019, Araujo et al. 2020, Desgarnier et al. 2022, Glaus et al. 2024).

Conclusion

While our understanding of coral reef ecosystem functioning remains in its infancy, this is yet another example of how preserving biodiversity could help safeguard important ecological mechanisms supporting

ecosystem health and associated benefits to society. We look forward to further studies on this topic, and advocate for increased protection of eagle rays and affiliate “ecological reef vacuums” as a potential nature based solution for regulating COTS populations and supporting coral reef management.

- Ajemian, M.J., Powers, S.P. and T.J.T. Murdoch. 2012. Estimating the potential impacts of large mesopredators on benthic resources: integrative assessment of spotted eagle ray foraging ecology in Bermuda. *PLOS ONE* 7: e40227. [CrossRef](#)
- Araujo, G., Legaspi, C., Matthews, K., Ponzo, A., Chin, A. and B.M. Manjaji-Matsumoto. 2020. Citizen science sheds light on the cryptic ornate eagle ray *Aetomylaeus vespertilio*. *Aquatic Conservation: Marine and Freshwater Ecosystems* 30: 2012–2018. [CrossRef](#)
- Bonham, B. and N.J. Silbiger. 2024. Exploring commensalism between rock wrasse (*Halichoeres semicinctus*) and round stingrays (*Urobatis halleri*) in Southern California. *Bulletin Southern California Academy of Sciences* 123: 53–59. [CrossRef](#)
- Broadhurst, M.K. and B.R. Cullis. 2020. Mitigating the discard mortality of non-target, threatened elasmobranchs in bather-protection gillnets. *Fisheries Research* 222: 105435. [CrossRef](#)
- Capapé, C. 1976. Etude du régime alimentaire de l'Aigle de mer, *Myliobatis aquila* (L., 1758) des côtes tunisiennes. *ICES Journal of Marine Science* 37: 29–35. [CrossRef](#)
- Cahill, B.V., Eckert, R.J., Bassos-Hull, K., Ostendorf, T.J., Voss, J.D., DeGroot, B.C. et al. 2023. Diet and Feeding Ecology of the Whitespotted Eagle Ray (*Aetobatus narinari*) from Florida Coastal Waters Revealed via DNA Barcoding. *Fishes* 8: 388. [CrossRef](#)
- Deaker, D.J., Agüera, A., Lin, H.A., Lawson, C., Budden, C., Dworjanyn, S.A., et al. 2020. The hidden army: corallivorous crown-of-thorns seastars can spend years as herbivorous juveniles. *Biology Letters* 16: 20190849. [CrossRef](#)
- Desgarnier, L., Mouillot, D., Vigliola, L., Chaumont, M. and L. Mannocci. 2022. Putting eagle rays on the map by coupling aerial video-surveys and deep learning. *Biological Conservation* 267: 109494. [CrossRef](#)
- Dulvy, N.K., Freckleton, R.P. and N.V. Polunin. 2004. Coral reef cascades and the indirect effects of predator removal by exploitation. *Ecology Letters* 7: 410–416. [CrossRef](#)
- Flowers, K.I., Heithaus, M.R. and Y.P. Papastamatiou. 2021. Buried in the sand: Uncovering the ecological roles and importance of rays. *Fish and Fisheries* 22: 105–127. [CrossRef](#)
- Glaus, K., Gordon, L., Vierus, T., Marosi, N.D., and H. Sykes. 2024. Rays in the shadows: Batoid diversity,

- occurrence, and conservation status in Fiji. *Biology* 13: 73. [CrossRef](#)
- Gray, A.E., Mulligan, T.J. and R.W. Hannah. 1997. Food habits, occurrence, and population structure of the bat ray, *Myliobatis californica*, in Humboldt Bay, California. *Environmental Biology of Fishes* 49: 227–238. [CrossRef](#)
- Harry, A.V., Tobin, A.J., Simpfendorfer, C.A., Welch, D.J., Mapleston, A., White, J., et al. 2011. Evaluating catch and mitigating risk in a multispecies, tropical, inshore shark fishery within the Great Barrier Reef World Heritage area. *Marine and Freshwater Research* 62: 710–721. [CrossRef](#)
- Heithaus, M.R., Frid, A., Wirsing, A.J. and B. Worm. 2008. Predicting ecological consequences of marine top predator declines. *Trends in Ecology & Evolution* 23: 202–210. [CrossRef](#)
- Heudier, M., Mouillot, D. and L. Mannocci. 2023. Assessing the effects of coral reef habitat and marine protected areas on threatened megafauna using aerial surveys. *Aquatic Conservation: Marine and Freshwater Ecosystems* 33: 286–297. [CrossRef](#)
- Hines, A.H., Whitlatch, R.B., Thrush, S.F., Hewitt, J.E., Cummings, V.J., Dayton, P., et al. 1997. Nonlinear foraging response of a large marine predator to benthic prey: eagle ray pits and bivalves in a New Zealand sandflat. *Journal of Experimental Marine Biology and Ecology* 216: 191–210. [CrossRef](#)
- Jardas, I., Šantić, M. and A. Pallaoro. 2004. Diet composition of the eagle ray, *Myliobatis aquila* (Chondrichthyes: Myliobatidae), in the Eastern Adriatic Sea. *Cybiurn*, 28: 372–374. [CrossRef](#)
- Kayal, M. 2024. Eagle ray feeding on a shallow coral reef in Kanaky New Caledonia. Aerial video footage, 2 min 45 sec. <https://youtu.be/sxdbiZJmX1Y>
- Kayal, M., Lenihan, H.S., Pau, C., Penin, L. and M. Adjeroud. 2011. Associational refuges among corals mediate impacts of a crown-of-thorns starfish *Acanthaster planci* outbreak. *Coral Reefs* 30: 827–837. [CrossRef](#)
- Kayal, M., Vercelloni, J., Lison de Loma, T., Bosserelle, P., Chancerelle, Y., Geoffroy, S., et al. 2012. Predator crown-of-thorns starfish (*Acanthaster planci*) outbreak, mass mortality of corals, and cascading effects on reef fish and benthic communities. *PLOS ONE* e47363. [CrossRef](#)
- Kroon, F.J., Barneche, D.R. and M.J. Emslie. 2021. Fish predators control outbreaks of Crown-of-Thorns Starfish. *Nature Communications* 12: 6986. [CrossRef](#)
- Lynch, A.M.J., Sutton, S.G. and C.A. Simpfendorfer. 2010. Implications of recreational fishing for elasmobranch conservation in the Great Barrier Reef Marine Park. *Aquatic Conservation: Marine and Freshwater Ecosystems* 20: 312–318. [CrossRef](#)
- McCook, L.J., Ayling, T., Cappo, M., Choat, J.H., Evans, R.D., De Freitas, D.M., et al. 2010. Adaptive management of the Great Barrier Reef: a globally significant demonstration of the benefits of networks of marine reserves. *Proceedings of the National Academy of Sciences*, 107: 18278–18285. [CrossRef](#)
- Mellin, C., MacNeil, A.M., Cheal, A.J., Emslie, M.J. and J.M. Caley. 2016. Marine protected areas increase resilience among coral reef communities. *Ecology Letters* 19: 629–637. [CrossRef](#)
- Myers, R.A., Baum, J.K., Shepherd, T.D., Powers, S.P. and C.H. Peterson. 2007. Cascading effects of the loss of apex predatory sharks from a coastal ocean. *Science* 315: 1846–1850. [CrossRef](#)
- Pratchett M.S. 2025. Establishing functionally important predators on crown-of-thorns starfish (*Acanthaster* spp.). *Ideas in Ecology & Evolution* 18: 32–34. [CrossRef](#)
- Sabino, J., Andrade, L.P., Sazima, I., Teresa, F.B., Floeter, S.R., Sazima, C. et al. 2016. Following fish feeding associations in marine and freshwater habitats. *Marine and Freshwater Research* 68: 381–387. [CrossRef](#)
- Schluessel, V., Bennett, M.B. and S.P. Collin. 2010. Diet and reproduction in the white-spotted eagle ray *Aetobatus narinari* from Queensland, Australia and the Penghu Islands, Taiwan. *Marine and Freshwater Research* 61: 1278–1289. [CrossRef](#)
- Serrano-Flores, F., Pérez-Jiménez, J.C., Méndez-Loeza, I., Bassos-Hull, K. and M.J. Ajemian. 2018. Comparison between the feeding habits of spotted eagle ray (*Aetobatus narinari*) and their potential prey in the southern Gulf of Mexico. *Journal of the Marine Biological Association of the United Kingdom* 99: 661–672. [CrossRef](#)
- Sporcic, M., Hobday, A., Bulman, C. and M. Fuller. 2018. Ecological risk assessment for the effects of fishing: Eastern tuna & billfish fishery: Longline Sub-fishery, data to 2015. Report for the Australian Fisheries Management Authority.
- Sumpton, W.D., Taylor, S.M., Gribble, N.A., McPherson, G. and T. Ham. 2011. Gear selectivity of large-mesh nets and drumlines used to catch sharks in the Queensland Shark Control Program. *African Journal of Marine Science* 33: 37–43. [CrossRef](#)
- Sweatman, H. 2008. No-take reserves protect coral reefs from predatory starfish. *Current Biology* 18: R598–9. [CrossRef](#)
- Tagliafico, A., Butcher, P.A., Colefax, A.P., Clark, G.F. and B.P. Kelaher. 2020. Variation in cownose ray *Rhinoptera neglecta* abundance and group size on the central east coast of Australia. *Journal of Fish Biology* 96: 427–433. [CrossRef](#)
- Vanhatalo, J., Hosack, G.R. and H. Sweatman. 2017. Spatiotemporal modelling of crown-of-thorns starfish outbreaks on the Great Barrier Reef to inform control strategies. *Journal of Applied Ecology* 54: 188–197. [CrossRef](#)

- Westcott, D.A., Fletcher, C.S., Kroon, F.J., Babcock, R.C., Plagányi, E.E., Pratchett, M.S. et al. 2020. Relative efficacy of three approaches to mitigate Crown-of-Thorns Starfish outbreaks on Australia's Great Barrier Reef. *Scientific Reports* 10: 12594. [CrossRef](#)
- Wilmes, J.C., Hoey, A.S. and M.S. Pratchett, 2020a. Contrasting size and fate of juvenile crown-of-thorns starfish linked to ontogenetic diet shifts. *Proceedings of the Royal Society B* 287: 20201052. [CrossRef](#)
- Wilmes, J.C., Schultz, D.J., Hoey, A.S., Messmer, V. and M.S. Pratchett. 2020b. Habitat associations of settlement-stage crown-of-thorns starfish on Australia's Great Barrier Reef. *Coral Reefs* 39: 1163–1174. [CrossRef](#)
- Yamaguchi, A., Kawahara, I. and S. Ito. 2005. Occurrence, growth and food of longheaded eagle ray, *Aetobatus flagellum*, in Ariake Sound, Kyushu, Japan. *Environmental Biology of Fishes* 74: 229–238. [CrossRef](#)